



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

There are, of course, many insects closely related to the Culicidæ, and at times some not even belonging to the dipteræ, sent in, but no effort is made to determine these, as they have no place in the problem on hand, though they are, many of them, of interest to the entomologist and may prove to be of interest to the pathologist.

Much of the country is still unexplored, so far as this research is concerned, the southern islands particularly having few collecting stations, and with the exception of the collections sent in by Major R. P. Ames, such as have been sent in have, partly owing to the change of surgeons, either been irregular or have not covered many consecutive months, so that other new species are to be expected and will undoubtedly be added to the list.

---

## TWO NEW HYMENOPTEROUS PARASITES.

BY WILLIAM H. ASHMEAD, M.A., D.Sc.

### ***Orgilus kearfotti*, sp. nov.**

♂. Length 3 mm. Honey-yellow, the head in front paler; disk of metathorax, the eyes and the stemmaticum black; flagellum dusky or brownish; a spot at apex of hind femora, the hind tibiæ, except a white annulus at base, and their tarsi, fuscous. Wings hyaline, the stigma and the veins brown.

Caldwell, N. J. One specimen, bred by Mr. W. D. Kearfott, July 2d, from *Recurvaria juniperella* Kearfott.

Type.—Cat. No. 6960, U. S. N. M.

### ***Protapanteles recurvariæ*, sp. nov.**

♀. Length 1.65 mm. Head and thorax black and shining, but sparsely punctate; the clypeus, the mandibles, scape of antennæ, the abdomen at sides and beneath, the second dorsal segment, and the legs, except a spot at apex of the hind femora, the apex of hind tibiæ and the hind tarsi except basally which are fuscous, honey-yellow. Wings hyaline, the stigma brown. The ovipositor is prominent clavate, black; the plate of the first segment is trapezoidal, sculptured, the rest of the abdomen smooth, impunctured.

Montclair, N. J. Two specimens, bred by Mr. W. D. Kearfott, June 2d and 7th from *Recurvaria piceaella* Kearfott. Another specimen was bred from *R. thujaella* Kearfott.

Type. Cat. No. 6961, U. S. N. M.